

Characteristics of supernova remnant G351.7+0.8 and a distance argument against its association with PSR J1721-3532

W. W. Tian^{1,2*}, M. Haverkorn^{3,4}, H. Y. Zhang¹

¹*National Astronomical Observatories, CAS, Beijing 100012, China*

²*Department of Physics & Astronomy, University of Calgary, Calgary, Alberta T2N 1N4, Canada*

³*Jansky Fellow, National Radio Astronomy Observatory; marijke@astro.berkeley.edu*

⁴*Astronomy Department, University of California-Berkeley, 601 Campbell Hall, Berkeley CA 94720, USA*

Accepted . Received ; in original form

ABSTRACT

New images of the supernova remnant (SNR) G351.7+0.8 are presented based on 21cm HI-line emission and continuum emission data from the Southern Galactic Plane Survey (SGPS). SNR G351.7+0.8 has a flux density of 8.4 ± 0.7 Jy at 1420 MHz. Its spectral index is 0.52 ± 0.25 ($S = \nu^{-\alpha}$) between 1420 MHz and 843 MHz, typical of adiabatically expanding shell-like remnants. HI observations show structures possibly associated with the SNR in the radial velocity range of -10 to -18 km/s, and suggest a distance of 13.2 kpc and a radius of 30.7 pc. The estimated Sedov age for G351.7+0.8 is less than 6.8×10^4 yrs. A young radio pulsar PSR J1721-3532 lies close to SNR G351.7+0.8 on the sky. The new distance and age of G351.7+0.8 and recent proper-motion measurements of the pulsar strongly argue against an association between SNR G351.7+0.8 and PSR J1721-3532. There is an unidentified, faint X-ray point source 1RXS J172055.3-353937 which is close to G351.7+0.8. This may be a neutron star potentially associated with G351.7+0.8.

Key words: supernova remnants: individual (G351.7+0.8) – pulsar: individual (PSR J1721-3532).

* E-mail: tww@iras.ucalgary.ca; tww@bao.ac.cn

1 INTRODUCTION

Supernova remnant (SNR) and pulsar (PSR) associations have been widely studied since the first radio pulsar was discovered in 1968. Associations are usually judged on criteria such as agreement in age and distance, angular proximity, and whether the transverse velocity derived for the pulsar is reasonable. However, an independent measure of the age and distance for the two objects is usually difficult. Often, the error in the measurement of the transverse velocity of a pulsar is comparable to the measurement itself, effectively rendering a non-detection. Another recently much-used method of confirming an association is searching for a pulsar wind nebula (PWN) in a SNR. If a pulsar is still inside its parent SNR, the high pressure of the surrounding plasma can confine the pulsar's relativistic wind generating synchrotron emission observable as a PWN. About 50 possible associations of SNR/PSR have been claimed in the literature (for a compilation see Tian & Leahy 2004, and additionally Tian & Leahy 2006), and a few of these associations have been contested (Nicastro et al. 1996; Stappers et al. 1999; Gaensler et al. 2001; Thorsett et al. 2002; Brisen et al. 2006).

G351.7+0.8 is a shell-type SNR first mapped by Whiteoak & Green (1996). It is a faint plateau of emission on the edge of a bright HII region. Gaensler & Johnston (1995) suggested a possible association with pulsar PSR J1721-3532, which was included in the SNR/PSR associations catalogue by Tian & Leahy (2004). In this paper, based on new radio observations of SNR G351.7+0.8 and recent proper-motion measurements of pulsar PSR J1721-3532, we give new physical parameters of G351.7+0.8 and argue against the association SNR G351.7+0.8/PSR J1721-3532.

2 OBSERVATIONS

The radio continuum and HI emission data sets come from the Southern Galactic Plane Survey (SGPS) which is described in detail by Haverkorn et al. (2006) and McClure-Griffiths et al. (2005). The SGPS images the HI line emission and 1.4 GHz continuum in the range $253^\circ < l < 357^\circ$, mostly in the fourth quadrant of our Galaxy, with the Australia Telescope Compact Array (ATCA) and the Parkes 64m single dish telescope. The continuum observations have a resolution of 100 arcsec and a sensitivity better than 1 mJy/beam. The HI data have an angular resolution of 2 arcmin, a rms sensitivity of ~ 1 K and the velocity resolution of the SGPS of 1 km s^{-1} . The continuum data are taken with the ATCA interferometer




Figure 1. The images of G351.7+0.8 at 1420 MHz from the SGPS with a beamsize of 100 arcsec, as shown in the lower right corner of the plot. The white cross and connected arrow show the position and proper motion direction of the pulsar PSR J1721-3532. The box in the map is used to estimate the flux density of G351.7+0.8.

only, which is insensitive to scales larger than 30 arcmin. Because the angular size of the SNR ($18 \text{ arcmin} \times 14 \text{ arcmin}$ in $l \times b$) is significantly smaller than this, all remnant emission is included in the continuum data.

3 RESULTS

3.1 Continuum Emission

Fig. 1 shows the image of G351.7+0.8 at 1420 MHz from the SGPS. The SGPS image has a lower resolution (100 arcsec) but higher sensitivity ($< 1 \text{ mJy/beam}$) than a previously published 843 MHz map (43 arcsec and $> 2 \text{ mJy/beam}$) from the Molonglo Observatory Synthesis Telescope (MOST) (Whiteoak & Green 1996), so the SGPS image shows more faint emission than the MOST map. The 1420 MHz and 843 MHz images have similar morphologies. The flux density of G351.7+0.8 is $8.4 \pm 0.7 \text{ Jy}$ at 1420 MHz and $11 \pm 1.1 \text{ Jy}$ at 843 MHz (Whiteoak & Green 1996), so that the spectral index between these two frequencies is 0.52 ± 0.25 ($S = v^{-\alpha}$). The Parkes 6 cm southern Galactic plane survey by Haynes, Caswell & Simons (1978) has such a low resolution ($4.1'$) that a flux estimate is very uncertain due to contamination from the neighboring HII region, but a rough estimate of the flux from G351.7+0.8 ($\sim 2 \text{ Jy}$) agrees well with the estimated spectral index.

3.2 Polarization

G351.7+0.8 does not show any polarization at 1.4 GHz. As the non-thermal emission from supernova remnants is intrinsically partially polarized, the lack of polarization indicates that the emission is depolarized. This depolarization is most likely due to varying polarization angle on scales smaller than the synthesized beam, caused by small-scale structure in electron density and/or magnetic field in or in front of the SNR. As SNRs at larger distances tend to exhibit structure on smaller angular scales, SNRs beyond a certain distance appear mostly depolarized. Comparison with many remnants with known dispersion measure (DM) indicates that G351.7+0.8 (DM= 979 pc cm⁻³ for a distance of 13.2 kpc, see section 4.2), is expected to be completely depolarized (Kothes & Landecker 2004), as observed.

3.3 HI Emission

We searched the SGPS radial velocity range from -247 km s⁻¹ to 165 km s⁻¹ for features in HI with similar morphology as G351.7+0.8. There is emission which is possibly coincident with the SNR only in the velocity range -10 to -18 km s⁻¹. Fig. 2 shows maps of HI emission associated with the SNR averaged over velocities in the ranges of [-11.54, -13.19], [-14.01, -15.66], and [-16.49, -18.14] km s⁻¹, flanked by two off-line channels. The HI spectrum over the whole velocity range, integrated over the whole remnant, is also shown. Superimposed on each map is a contour of 1420 MHz continuum at 58 mJy/beam, chosen to denote the position and extent of the SNR. Fig. 3 shows the column density map of the HI integrated over channels from -10 to -18 km/s. Continuum subtraction has been performed in all plots.

4 DISCUSSION

4.1 The Distance and age

Using a flat Galactic rotation velocity $V_R=V_0=200$ km s⁻¹ and $R_0=8.0$ kpc, we obtain the SNR's distance $d = 2.7 \pm 0.6$ kpc or $d = 13.2 \pm 0.5$ kpc. The angular size of G351.7+0.8 is 18' by 14' (average diameter 16'), so this yields a radius of about 6 pc ($d = 2.7$ kpc) or 30.7 pc ($d = 13.2$ kpc) for the SNR. From the column density of the HI associated with G351.7+0.8 (see Fig. 3), we estimate an approximate N_H of $1 - 5 \times 10^{20}$ cm⁻², so the approximate density $n_0=N_H/(2R)$ is about 7.8 cm⁻³ ($d = 2.7$ kpc) or 1.6 cm⁻³ ($d = 13.2$ kpc). An ISM density of 7.8 cm⁻³ is so high that we should have seen HI piled up in a clear shell around the SNR,




Figure 2. HI emission in the field centered on G351.7+0.8. The first five frames (left to right, top to bottom) show HI emission at -8.24 km s $^{-1}$, averaged over velocity ranges $[-11.54, -13.19]$, $[-14.01, -15.66]$, and $[-16.49, -18.14]$ km s $^{-1}$, respectively. A contour of the 1420 MHz continuum at 58 mJy/beam overlaid on each map. The bottom right plot shows the HI spectrum over the complete velocity range integrated over the whole remnant. The inlay shows the velocity region in which the HI is morphologically similar to the continuum emission from the remnant.

which is not seen in Fig. 2. The absence of such a shell means G351.7+0.8 is most likely at the farther distance of 13.2 kpc.

Applying a Sedov model (Cox 1972), for a typical explosion energy of $E = 0.5 \times 10^{51}$ erg ($\epsilon_0 = E/(0.75 \times 10^{51} \text{ erg}) = 2/3$), yields an age of 1.4×10^5 yr for $d = 13.2$ kpc.

Because the Sedov phase only lasts for several 10,000 years, the SNR would have com-




Figure 3. Column density of the HI in the field centered on G351.7+0.8, in 10^{20} atoms cm^{-2} . The contour is same as in Fig. 2.

pletely cooled by that time for $n_0 = 1.6 \text{ cm}^{-3}$. The shock radius when the SNR has cooled is $R_s^{(c)} = 24.3(\epsilon_0/n_0)^{5/17} n_0^{-2/17} \text{ pc}$. Because the remnant would show a thin, bright shell if it was older than a cooling time, contrary to observations, we require that the SNR age is smaller than the cooling time, i.e. $R \leq R_s^{(c)}$. Thus we find an upper limit for n_0 of 0.4 cm^{-3} . This is consistent with a HI column density of about $0.8 \times 10^{20} \text{ cm}^{-2}$, roughly consistent with Fig. 3. The revised age for G351.7+0.8 with $n_0 < 0.4 \text{ cm}^{-3}$ is less than $6.8 \times 10^4 \text{ yr}$.

4.2 SNR G351.7+0.8 and PSR J1721-3532

Pulsar J1721-3532 ($l=351.69^\circ$, $b=0.67^\circ$) is situated at a slightly lower latitude than SNR G351.7+0.8 and obscured by a small HII region. It has a characteristic age of 178 kyr (Taylor et al. 1993) and a DM distance of 5.6 kpc based on the electron density model of Cordes & Lazio (2002). The pulsar is shown by the white cross in Fig. 1, where the arrow denotes its proper motion direction. The angular separation between the pulsar and the centre of the SNR is $\sim 10'$. Weisberg et al. (1995) gave a lower limit to the distance of the pulsar of $4.8 \pm 0.3 \text{ kpc}$ from HI absorption spectra. Although no distance and age are available for SNR G351.7+0.8, Gaensler and Johnston (1995) mentioned a possible association of SNR G351.7+0.8 with PSR J1721-3532 based on their β parameter method, i.e., their calculations showed a pulsar this old could have traveled outside the SNR in its lifetime. However, the SNR's parameters given in the paper make this association unlikely.

The Cordes & Lazio (2002) electron density model yields $\text{DM} = 979 \text{ pc cm}^{-3}$ for the SNR at a distance of 13.2 kpc. The estimated distance error in the DM values is about 5 per cent

in this direction (Cordes & Lazio 2002). The pulsar J1721-3532 has a DM of $496 \pm 4 \text{ pc cm}^{-3}$, which is widely inconsistent with the DM estimate for the SNR.

4.3 Timing Measurements for PSR J1721-3532

A pulsar proper motion measurement is a good way to support SNR/PSR associations or disprove candidate associations. Recent timing measurements (Zou et al. 2005) gave PSR J1721-3532's proper motion $7200 \pm 2400 \text{ km s}^{-1}$ at a distance of 5.6 kpc ($\mu_\alpha = -43 \pm 24 \text{ mas yr}^{-1}$, $\mu_\delta = -270 \pm 93 \text{ mas yr}^{-1}$). Zou et al.'s (2005) proper motion measurements are based upon comparing two position estimates (probably including a significant underestimate of position uncertainties) and cite large errors which are probably related with a lot of timing noise from the pulsar, adding to the position uncertainty. However, the proper motion direction is still more or less similar if the true values of the velocity are within 2σ from the cited value, which adds weak evidence against the association PSR J1721-3532/SNR G351.7+0.8 (see Fig. 1).

4.4 The Rosat point source 1RXS J172055.3-353937

G351.7+0.8 is an old SNR, its soft X-ray emission is not expected to be bright in a lower density environment, and should have been completely absorbed at a distance of 13.2 kpc. This is consistent with lack of detection within G351.7+0.8 in the Rosat All-sky Survey. There is an unidentified, faint X-ray point source, listed by Voges et al. (1999) as 1RXS J172055.3-353937. It is centred on $l=351.52^\circ$, $b=0.71^\circ$, and lies close to the southeastern edge of G351.7+0.8. This X-ray source has a total of 6 counts in the Rosat All-Sky survey, not enough to deduce any spectral information. The source does not seem to have an optical counterpart. It may coincide with a hard X-ray source in Fig. 2 of Ezoe et al (2006), possibly indicating a large distance at which soft X-rays have been absorbed. Obviously, deeper X-ray observations are needed. If it is confirmed to be a neutron star later, it is reasonable to associate G351.7+0.8 with 1RXS J172055.3-353937.

5 CONCLUSION

We present new images of the SNR G351.7+0.8 from the 1.4 GHz radio continuum and HI emission dataset from the SGPS. Its continuum flux density at 1.4 GHz is $8.4 \pm 0.7 \text{ Jy}$ and its spectral index is 0.52 ± 0.25 . Based on associated HI emission, the remnant's

distance is estimated to be 13.2 ± 0.5 kpc. No polarization is detected, consistent with the expectation of complete depolarization at this distance towards the Galactic centre. Based on the assumption that the remnant must be younger than a cooling time, its age is younger than 6.8×10^4 yrs, and the ambient ISM density is $n_0 < 0.4 \text{ cm}^{-3}$. Discrepancies in distance and dispersion measure, and a tentative proper motion measurement of pulsar J1721-3532, make an association between the two objects SNR G351.7+0.8 and PSR J1721-3532 unlikely. A faint unidentified X-ray point source, 1RXS J172055.3-353937, is a potential candidate to be associated with G351.7+0.8.

ACKNOWLEDGMENTS

We thank the referee for his/her comments that have improved the paper. TWW acknowledges supports from the Natural Sciences Foundation of China and the Natural Sciences and Engineering Research Council of Canada. MH acknowledges support from the National Radio Astronomy Observatory (NRAO), which is operated by Associated Universities Inc., under cooperative agreement with the National Science Foundation. We thank Dr. Kothes for useful discussions, and thank the SGPS team for the data.

REFERENCES

- Briskin W.F., Carrillo-Barragan M., Kurtz S., Finley J.P., 2006, *ApJ*, 652, 554
 Cordes J.M., Lazio T.J.W., 2002, preprint (astro-ph/0207156)
 Cox D., 1972, *ApJ*, 178, 159
 Ezoe Y., Kokubun M., Makishima K., Sekimoto Y., Matsuzaki K., 2006, *ApJ*, 638, 860
 Gaensler B.M., Slane P.O., Gotthelf E.V., Vasisht G., 2001, *ApJ*, 559, 963
 Gaensler B.M., Johnston S., 1995, *MNRAS*, 275, 73
 Haverkorn M., Gaensler B.M., McClure-Griffiths N.M., Dickey J.M., Green A.J., 2006, *ApJS*, accepted
 Haynes R.F. Caswell J.L., Simons L.W.J., 1978, *AuJPA*, 45, 1
 Kothes R., Landecker T.L., 2004, in "The Magnetized Interstellar Medium", Eds. B. Uyaniker, W. Reich, R. Wielebinski, Copernicus GmbH, p. 33
 McClure-Griffiths N.M., Dickey John M., Gaensler B.M., Green A.J., Haverkorn M., Strasser S., 2005, *ApJS*, 158, 178
 Nicastro L., Johnston S., Koribalski B., 1996, *A&A*, 306, 49

- Stappers B.W., Gaensler B.M., Johnston S., 1999, MNRAS, 308, 609
- Taylor J.H., Manchester R.N., Lyne A.G., 1993, ApJS, 88, 529
- Thorsett S.E., Bricken W.F., Goss W.M., 2002, ApJ, 573, L111
- Tian W.W., Leahy D.A., 2006, A&A, 455, 1053
- Tian W.W., Leahy D.A., 2004, Progress in Astronomy, 22, 308
- Voges W., Aschenbach B., Boller T.H., 1999, A&A, 349, 389
- Weisberg J.M., Siegel M.H., Frail D.A., Johnston S., 1995, ApJ, 447, 204
- Whiteoak J.B.Z., Green A.J., 1996, A&AS, 118, 329
- Zou W.Z., Hobbs G., Wang N., Manchester R.N., Wu X.J., Wang H.X., 2005, MNRAS, 362, 1189